Screening the Antibacterial Activities of *Streptomyces* Extracts against Phytopathogens *Xanthomonas oryzae* pathovar *oryzae*, *Xanthomonas campestris* pathovar *vesicatoria*, and *Pectobacterium carotovorum* pathovar *carotovorum*

Seung-Hwan Kim · Jinhua Cheng · Seung Hwan Yang · Joo-Won Suh · Eun-Sung Song · Lin-Woo Kang* · Jeong-Gu Kim*

Received: 7 May 2015 / Accepted: 7 July 2015 / Published Online: 30 September 2015
© The Korean Society for Applied Biological Chemistry 2015

Abstract *Xanthomonas oryzae* pv. *oryzae* (*Xoo*), *X. campestris* pv. *vesicatoria* (*Xcv*), and *Pectobacterium carotovorum* pv. *carotovorum* (*Pcc*) are the causative agents of bacterial blight in rice, bacterial spot in pepper, and bacterial soft rot in carrot and cabbage, respectively. To isolate novel microbial extracts with antimicrobial activities against these bacteria, approximately 5,300 different *Streptomyces* extracts were prepared and tested. Microbial cultures from various *Streptomyces* strains isolated from the Jeju Island, Baekam, Mankyong river, Jiri mountain etc. in Korea were extracted into three different factions -secreted hydrophobic, secreted hydrophilic, and mycelia- using ethyl acetate, water, and methanol. Initially, 34, 29, and 10 extracts were selected as having antibacterial activities against *Xoo*, *Xcv*, and *Pcc*, respectively. Extracts 1169G4, 1172E9, and 1172E10 had the

highest growth inhibition activities against both *Xoo* and *Xcv*, and extracts 1151H7 and 1152H7 showed the highest growth inhibition activities against *Pcc*.

Keywords antimicrobial activity · *Pectobacterium carotovorum* pathovar *carotovorum* · *Streptomyces* · *Xanthomonas campestris* pathovar *vesicatoria* · *Xanthomonas oryzae* pathovar *oryzae*

Introduction

Filamentous bacteria belonging to the genus *Streptomyces* produce over 50% of the antibiotics discovered to date (Taddei et al., 2006; Jayapal et al., 2007). *Streptomyces* bacteria and other Actinomycetes produce a diverse array of antibiotics, including aminoglycosides, anthracyclins, glycopeptides, β -lactams, macrolides, nucleosides, peptides, polyenes, polyethers, and tetracyclines (Mellouli et al., 2003).

Rice, pepper, and cabbage are the first, second, and third most cultivated crops, respectively, in Korea. *Xanthomonas oryzae* pv. *oryzae* (*Xoo*) and *X. campestris* pv. *vesicatoria* (*Xcv*) are the causal agents of bacterial blight in rice and bacterial spot in pepper, respectively (Yang et al., 2005; Adhikari et al., 2010). *Pectobacterium carotovorum* pv. *carotovorum* (*Pcc*) causes soft rot in a wide range of plants, including cabbage (Mole et al., 2010). Because these bacteria cause severe crop damage and reduce yield, prevention and treatment of these diseases have been the focus of intense research in both the academy and industry (Khush, 1997). In 2005, the whole genomes of *Xoo* and *Xcv* were sequenced (Frank Thieme, 2005; Lee et al., 2005). The emergence of resistance against *Pcc* was reported in 2007 (T. Luzzatto, 2007).

Chemical pesticides are routinely used to control plant pathogens in agricultural fields. However, recently, indiscriminate usage of

S.-H. Kim · E.-S. Song · J.-G. Kim
Genomics Division, National Academy of Agricultural Science (NAAS), Rural Development Administration (RDA), Jeonju 560-500, Republic of Korea

J. Cheng · J.-W. Suh
Division of Bioscience and Bioinformatics, College of Natural Science, Myongji University, Yongin 449-728, Republic of Korea

S. H. Yang
Interdisciplinary Program of Biomodulation, Myongji University, Yongin 449-728, Republic of Korea

L.-W. Kang
Department of Biological Sciences, Konkuk University, Hwayang dong, Gwangjin-gu, Seoul 143-701, Republic of Korea

*Corresponding authors (L.-W. Kang: lkang@konkuk.ac.kr; J.-G. Kim: jkim5aug@korea.kr)

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

chemical pesticides has raised many concerns, including environmental toxicity (Vivek K. Bajpai, 2010) and the possibility of generating antibiotic-resistant microbes. Therefore, biopesticides made from secondary metabolites from microbes or plants are being preferred over chemical pesticides.

The main objective of this study was to isolate novel antimicrobial *Streptomyces* extracts against the phytopathogens mentioned. *Streptomyces* species are ubiquitous in the soil, and play an important ecological role in the turnover of organic material. Furthermore, the diverse secondary metabolites produced by *Streptomyces* species are good sources of novel bioactive compounds. To date, more than half of the world's antibiotics (over 50 different antibiotics) have been isolated from *Streptomyces*. In this study, we prepared and screened 5,328 *Streptomyces* extracts, and tested their antimicrobial activities against *Xoo*, *Xcv*, and *Pcc*.

Materials and Methods

Strains and Culture Conditions. *X. oryzae* pv. *oryzae* KACC10331 (*Xoo*), *X. campestris* pv. *vesicatoria* KACC11127 (*Xcv*) and *P. carotovorum* pv. *carotovorum* KACC10227 (*Pcc*) were provided by the Korea Agriculture Culture Collection (KACC). *Xoo*, *Xcv*, and *Pcc* were grown in YPD (1.0% yeast extract, 2.0% peptone, and 2.0% dextrose) rich media at 28°C. *Streptomyces* strains, collected from various regions of the Republic of Korea, including Jeju island, Baekam, Mankyong river, Jiri mountain etc. (S-Table 1), were maintained at the Extract Collection of Useful Microorganisms library at the Myoungji University, and grown at 28°C on GSS medium (1.0% soluble starch, 2.0% glucose, 0.1% beef extract, 0.4% yeast extract, 0.2% NaCl, 0.025% K₂HPO₄, and 0.2% CaCO₃ at pH 7.2), Bennett's medium (1.0% glucose, 0.1% yeast extract, 0.2% bacto-peptone, and 0.1% beef extract), or NYC medium (2.5% dextrin, 1.2% dry yeast, 2.0% corn starch liqueur, 0.1% NaBr, and 0.1% CoCl₂ at pH 7.0).

Preparation of the *Streptomyces* extracts. For preparation of *Streptomyces* extracts library, one gram of soil were suspended in 10 milliliter of 0.85% NaCl solution and serially diluted. The

diluted soil suspension were spread on Humic acid- vitamin (HV) agar medium (0.1% humic acid, 0.05% Na₂HPO₄, 0.071% KCl, 0.005% MgSO₄·7H₂O, 0.001% FeSO₄·7H₂O, 0.002% CaCO₃, 0.2% B-Vitamin solution, 1.8% agar at pH 7.2) and incubated at 28°C for 14 days. The colony with representative actinomycetes were selected and identified by 16S rDNA sequence. Then *Streptomyces* strains were cultured in three different media: GSS medium for nutrient-rich condition, Bennett's medium for moderate-nutrition condition, and NYC medium for nutrient-poor condition. Three extract fractions were prepared from each *Streptomyces* culture broth. 30 mL of the culture broth were centrifuged at 5,000×g for 10 min. Mycelia were collected as a pellet, and extracted with 3 mL of methanol. This fraction was called "mycelia isolated from supernatant (MIS)". After pelleting the mycelia, the supernatant was further extracted with 30 mL of ethyl acetate and dried. The resulting dried extract was dissolved in 3 mL of methanol. This was called the "secreted hydrophobic (SHB) fraction". The remaining ethyl acetate-insoluble fraction was called the "secreted hydrophilic (SHL) fraction". Thirty microliter of each fraction was distributed into 96-well plate and dried completely before further use. The dry weights were measured to enable accurate calculation of GI50 (dose required to cause 50% growth inhibition).

Naming the *Streptomyces* extracts. All extracts were represented as "0000X0." The four digits, preceding the alphabet "X," refer to the plate number, which is named according to the region from where the sample was collected. The alphabet X refers the row in which the sample was present (numbered from top to bottom) and the last digit represents the column (numbered from left to right). For instance, sample 1172H2 is located in the second column from the left and at the bottom row of the 96-well plate numbered 1172. In addition, the last digit of the extract's name represents the extract-type: 2, 5, and 8 represent SHL; 3, 6, and 9 represent MIS; and 4, 7, and 10 represent SHB.

Antimicrobial activity test. The antimicrobial activities of the extracts were estimated by the growth inhibition test using a single dose in two steps. In the first round of screening, all the extracts were stored in columns 2–10 of the 96-well plate. Columns 1, 11, and 12 contained the control (medium only), bacterial culture, and

Table 1 Antibacterial extracts screened in the first and second round against *Xoo*, *Xcv*, and *Pcc*

	<i>Xoo</i>			<i>Xcv</i>		<i>Pcc</i>	
	SHL	1119D8, 1117E2		1128D2, 1128B5	1128B2	1158H5, 1158G2	
1 st	SHB	1176A4, 1175D7, 1175A4, 1173E4, 1172E10, 1169H4, 1169G4, 1159B10, 1149F4, 1139G10, 1136G7, 1124A7, 1120F10, 1119F10, 1119C10, 1117E4		1175A4, 1169G4, 1169E10, 1144H7, 1144A7, 1141H7, 1141A10, 1133H10, 1132H7, 1131H4, 1128H4, 1128C4, 1128B4, 1121F10	1120F10, 1172E10	1153E7, 1152H7, 1151H7, 1158G7	
	MIS	1175A3, 1173E3, 1172E9, 1169G3, 1164E6, 1161F3, 1161E3, 1160D3, 1159B9, 1158A6, 1149F3, 1144B6, 1135H3, 1120F9, 1119C6, 1117E3		1172E9, 1152H3, 1144A6, 1141H6, 1132H6, 1131H3, 1128H6, 1128D3, 1128C6, 1128A3		1158H6, 1158G6, 1157A6, 1155E9	
	SHL	-		-		-	
2 nd	SHB	1172E10, 1169G4		1172E10, 1169G4		1152H7, 1151H7	
	MIS	1172E9		1172E9		-	

bacterial culture with 6% dimethylsulfoxide (DMSO), respectively. The *Streptomyces* extracts were dissolved in 40 μ L of 50% DMSO, and further diluted in 100 μ L of YPD medium. *Xoo*, *Xcv*, and *Pcc* were grown to the early exponential growth stage (absorbance at 600 nm [A_{600}]=0.2), and 160 μ L of each bacterial culture was dispensed into separate 96-well micro-plates (Nunc $\text{\textregistered}$, US). Then, 40 μ L of previously diluted *Streptomyces* extract was added to each well. The cultures were grown at 28°C with continuous shaking at 700 rpm. The A_{600} values of the wells in the culture plates were measured every 4 h for 20 h using a multimode micro-plate reader (Infinite $^{\text{®}}$ M200 PRO, TECAN, Switzerland). To measure the antibacterial activity of each extract, we calculated the growth inhibition rate (GIR) at 20 h for *Xoo* and *Xcv* and at 12 h for *Pcc* using the following formula:

$$GIR = (1[CB]/[AB]) \times 100$$

where A was the A_{600} value of the bacterial culture with 6% DMSO; $B=0.2$, which was the A_{600} value of the bacterial culture when the extract was added; and C was the A_{600} value of the bacterial culture after treatment with the extract. For the second growth inhibition screen, the extracts selected from the first assay were tested three times using the same procedure.

The amount of growth inhibition induced by each extract was measured at five different dosages (1 , 10^{-1} , 10^{-2} , 10^{-3} , and 10^{-4}

Fig. 1 A bar pie chart depicting the results of the first round of screening of the *Streptomyces* extracts for antimicrobial activity against (A) *Xanthomonas oryzae* pv. *oryzae* (*Xoo*), (B) *X. campestris* pv. *vesicatoria* (*Xcv*), and (C) *Pectobacterium carotovorum* pv. *carotovorum* (*Pcc*). The bar lists screened antimicrobial extracts that showed more than 50% growth inhibition. The percentages of screened extracts are indicated. Extracts showing growth inhibition of 51–80 and 81–100% are indicated separately in the enlarged bar. The *Streptomyces* extracts were prepared as described in the Materials and Methods.

dilutions) and compared with the controls. From this, the GI50 was calculated. All the tests were carried out in triplicate.

Fig. 2 First round of screening for antibacterial extracts against (A) *Xoo*, (B) *Xcv*, and (C) *Pcc*. The bar depicts the relative growth rate of (A) *Xoo*, (B) *Xcv*, and (C) *Pcc* upon treatment with each screened extract, compared with the control (untreated bacterial culture). The relative growth for control was defined 1.0. Extracts are labeled as “0000X0,” as described in the Materials and Methods. The first four numbers and the last two characters are labeled separately.

Fig. 3 The growth curve of (A) *Xoo*, (B) *Xcv*, and (C) *Pcc* upon treatment with the screened antibacterial extracts in the second round of growth-inhibition screening. The extracts were added at the early exponential growth stage (absorbance at 600 nm [A_{600}]=0.2, time=0) of each bacterial culture, and bacterial growth was monitored for an additional 12–15 h by measuring A_{600} . Untreated bacterial culture was used as the control sample.

Results

Preparation of extracts from *Streptomyces*. *Streptomyces* strains produce diverse physiological secondary metabolites depending on the culture conditions. To generate various secondary metabolites from each *Streptomyces* strain, three culture media with different nutritional levels were used to grow the *Streptomyces* cells. In addition, three extract fractions were prepared for each *Streptomyces* culture. The MIS was separated from the supernatant; SHBs were extracted from the supernatant fraction using ethyl acetate; and SHLs were obtained from the residual water layer. In total, 5,328 extracts were prepared and screened for their antimicrobial activities against the three plant pathogens, *Xoo*, *Xcv*, and *Pcc*.

First round of screening for antimicrobial activity. Extracts showing more than 50% cell growth inhibition activity were considered antimicrobial extracts; extracts 34, 29, and 10 were selected because they exhibited antimicrobial activity against *Xoo*, *Xcv*, and *Pcc*, respectively. These three extracts constitute 0.64, 0.53, and 0.19% of all the extracts tested in the study (Fig. 1 and Table 1).

Extract 1172H9, the most effective antimicrobial extract against *Xoo*, inhibited 90% of *Xoo* cell growth. Of the top 34 extracts against *Xoo*, two were SHLs, 16 were MISs, and 16 were SHBs (Fig. 2A). Extract 1131H3 exhibited approximately 100% growth-inhibition effect against *Xcv*, and was the most effective antimicrobial against *Xcv*. Of the top 29 extracts exhibiting growth-inhibition effect against *Xcv*, three were SHLs, 10 were MISs, and 16 were SHBs (Fig. 2B). Extract 1152H7 showed 90% growth-inhibition activity against *Pcc*, and was the most effective antimicrobial extract against *Pcc*. Of the top 10 extracts showing growth-inhibition effect against *Pcc*, two were SHLs, four were MISs, and

the remaining four were SHBs (Fig. 2C).

Second round of screening for antimicrobial activity. Extracts showing high growth-inhibition activity in the first round of screening were tested three times to confirm their antimicrobial activities. 1169G4 (SHB), 1172E9 (MIS), and 1172E10 (SHB) had the highest growth-inhibition activities against both *Xoo* and *Xcv* (Fig. 3A and 3B). 1172E9 (MIS) and 1172E10 (SHB) are from the same *Streptomyces* strain from Jeju island. 1169G4 (SHB) is also from Jeju island (S-Table 1). 1152H7 (SHB) and 1151H7 (SHB) showed the highest growth-inhibition activity against *Pcc* (Fig. 3C). 1152H7 (SHB) and 1151H7 (SHB) are from Baekam, Gyoungsangbukdo (S-Table 1). 1152H7 (SHB) showed slightly higher antibacterial activity than 1151H7 (SHB). In total, five different extracts -1169G4 (SHB), 1172E9 (MIS), 1172E10 (SHB), 1152H7 (SHB), and 1151H7 (SHB)- were found to possess antibacterial activities against *Xoo*, *Xcv*, and *Pcc*. Interestingly, all the extracts, except 1172E9 (MIS), were SHBs.

The GI50s of the five antimicrobial extracts were measured by serial dilution of the extracts (Table 2). 1172E9 (MIS) and 1172E10 (SHB) showed the highest antibacterial activities against *Xoo* and *Xcv* (Fig. 4 and 5). In particular, 1172E10 (SHB) showed a GI50 value of less than $\mu\text{g}\cdot\text{mL}^{-1}$. 1152H7 (SHB) had a slightly

Table 2 Doses of screened antibacterial extracts required to cause 50% inhibition (GI50s) (Unit: $\mu\text{g}\cdot\text{mL}^{-1}$)

	<i>Xoo</i>	<i>Xcv</i>	<i>Pcc</i>
1169G4	4.1	1.7	-
1172E9	<1.7	<1.7	-
1172E10	0.3	0.4	-
1151H7	-	-	2.2
1152H7	-	-	0.5

Fig. 4 The growth curves of *Xoo* treated with (A) 1169G4, (B) 1172E9, and (C) 1172E10.

Fig. 5 The growth curves of *Xcv* treated with (A) 1169G4, (B) 1172E9, and (C) 1172E10.

higher antibacterial activity than 1151H7 (SHB) against *Pcc* (Fig. 6). The antimicrobial activities of all five extracts were less than $\mu\text{g}\cdot\text{mL}^{-1}$. 1172E10 (SHB: MJM strain No. 10146GE) had the highest antimicrobial activity against *Xoo* and *Xcv* with GI50s of 0.3 and 0.4 $\mu\text{g}\cdot\text{mL}^{-1}$, respectively. 1152H7 (SHB: MJM strain No. 9973DE) had the highest antimicrobial activity against *Pcc*, with a GI50 of 0.5 $\mu\text{g}\cdot\text{mL}^{-1}$.

Discussion

Antibiotics are secondary metabolites that inhibit the growth of other microbial organisms. Growing incidences antibiotic-resistance among deadly pathogens has increased the need to discover novel antibiotics. Over the last few decades, new technologies for drug discovery have been developed, such as high throughput screening

Fig. 6 The growth curves of *Pcc* treated with (A) 1151H7 and (B) 1152H7.

and combinatorial chemistry. Moreover, sequencing of pathogen genomes has led to the identification of new drug targets, and many chemical compounds have been synthesized by combinatorial chemistry and screened using cell-free assay systems and high-throughput screening systems. However, most screened compounds do not penetrate the pathogen to reach the target, or are easily degraded *in vivo*. Despite these technological advances, only five new classes of antibiotics have been introduced for human use (Talbot et al., 2006). Therefore, many studies are now focusing on screening secondary metabolites from microbes, especially *Streptomyces*, to identify novel antimicrobials. *Streptomyces* bacteria produce numerous secondary metabolites, and approximately 55% of currently used antibiotics are produced by the genus *Streptomyces*.

Five *Streptomyces* extracts showing high antimicrobial activity against *Xoo*, *Xcv*, and *Pcc* were screened. Of the five extracts, four (1152H7, 1151H7, 1169G4, and 1172E10) were secreted and hydrophobic, and only one extract (1172E9) was non-secreted. The secretory and hydrophobic properties of the extracts aid in the penetration of bacterial cell walls (Nikaido H, 1976; Savage et al., 2002). In particular, 1169G4, 1172E9, and 1172E10 were effective against both *Xcv* and *Xoo*. The genus *Xanthomonas* belongs to Proteobacteria, and it causes a variety of diseases in numerous crops. The extracts identified in this study may have applications as a novel antimicrobial drug against *Xanthomonas*, including *Xoo* and *Xcv*.

Acknowledgments This work was supported by grants from the RDA Next-Generation BioGreen 21 Program (No. PJ01132401).

References

- Adhikari SR, Supakunkunti S, and Khan MM (2010) Kala azar in Nepal: estimating the effects of socioeconomic factors on disease incidence. *Kathmandu Univ Med J (KUMJ)* **8**, 73–9.
- Frank Thieme RK, Thomas Bekel, Carolin Berger, Jens Boch, Daniela Büttner, Camila Caldana et al. (2005) Insights into Genome Plasticity and Pathogenicity of the Plant Pathogenic Bacterium *Xanthomonas campestris* pv. *vesicatoria* Revealed by the Complete Genome Sequence. *Journal of Bacteriology* **187**, 7254–66.
- Jayapal KP, Lian W, Glod F, Sherman DH, and Hu WS (2007) Comparative genomic hybridizations reveal absence of large *Streptomyces coelicolor* genomic islands in *Streptomyces lividans*. *Bmc Genomics* **8**, 229.
- N. Huang, E. R. Angeles, J. Domingo, G. Magpantay, S. Singh, G. Zhang et al. (1997) Pyramiding of bacterial blight resistance genes in rice: marker-assisted selection using RFLP and PCR. *TAG Theoretical and Applied Genetics* **95**, 313–20.
- Lee BM, Park YJ, Park DS, Kang HW, Kim JG, Song ES et al. (2005) The genome sequence of *Xanthomonas oryzae* pathovar *oryzae* KACC10331, the bacterial blight pathogen of rice. *Nucleic Acids Res* **33**, 577–86.
- Mellouli L, Ben Ameer-Mehdi R, Sioud S, Salem M, and Bejar S (2003) Isolation, purification and partial characterization of antibacterial activities produced by a newly isolated *Streptomyces* sp US24 strain. *Research in Microbiology* **154**, 345–52.
- Mole B, Habibi S, Dangel JL, and Grant SR (2010) Gluconate metabolism is required for virulence of the soft-rot pathogen *Pectobacterium carotovorum*. *Mol Plant Microbe Interact* **23**, 1335–44.
- Nikaido H (1976) Outer membrane of *Salmonella typhimurium*. Transmembrane diffusion of some hydrophobic substances. *Biochimica et Biophysica Acta* **433**, 118–32.
- Savage PB, Li C, Taotafa U, Ding B, and Guan Q (2002) Antibacterial properties of cationic steroid antibiotics. *FEMS Microbiol Lett* **217**, 1–7.
- T. Luzzatto MY Yishay, A. Lipsky, A. Ion, E. Belasov, and I. Yedidia (2007) Efficient, long-lasting resistance against the soft rot bacterium *Pectobacterium carotovorum* in calla lily provided by the plant activator methyl jasmonate. *Plant Pathology* **56**, 692–701.
- Taddei A, Rodriguez MJ, Marquez-Vilchez E, and Castelli C (2006) Isolation and identification of *Streptomyces* spp. from Venezuelan soils: Morphological and biochemical studies. I. *Microbiological Research* **161**, 222–31.
- Talbot GH, Bradley J, Edwards JE, Jr., Gilbert D, Scheld M, and Bartlett JG (2006) Bad bugs need drugs: an update on the development pipeline from the Antimicrobial Availability Task Force of the Infectious Diseases Society of America. *Clin Infect Dis* **42**, 657–68.
- Vivek K. Bajpai NTD, Hwa-Jin Suh, and Sun Chul Kang (2010) Antibacterial Activity of Essential Oil and Extracts of *Cleistanthus operculatus* Buds Against the Bacteria of *Xanthomonas* spp. *Journal of the American Oil Chemists' Society* **87**, 1341–9.
- Yang W, Sacks EJ, Lewis Ivey ML, Miller SA, and Francis DM (2005) Resistance in *Lycopersicon esculentum* Intrasppecific Crosses to Race T1 Strains of *Xanthomonas campestris* pv. *vesicatoria* Causing Bacterial Spot of Tomato. *Phytopathology* **95**, 519–27.