Structure elucidation of 11-epiterpestacin glycoside (11-ETG) isolated from *Bipolaris sorokiniana* NSDR-011

Chi-Hwan Lim¹ · Youichirou Nihashi²

Received: 8 February 2018 / Accepted: 26 February 2018 / Published Online: 31 March 2018
© The Korean Society for Applied Biological Chemistry 2018

Abstract An α -D-glucoside of sesterterpene, 11-epiterpestacin, was isolated from the culture of a filamentous fungus *Bipolaris sorokiniana* NSDR-011. The structure was elucidated by chemical studies and spectroscopic methods including NMR and ESI-MS. 11-ETG (**1**) named arbitrarily did not inhibit the root growth of Italian ryegrass seedlings even at the level of 200 ppm, while its aglycone 11-ET (**2**) completely inhibited root growth at level of 100 ppm.

Keywords *Bipolaris sorokiniana* · 11-epiterpestacin (11-ET) · 11-epiterpestacin glucoside (11-ETG) · Phytotoxin · Sesterterpene

Introduction

Plant pathogenic fungi are well known to produce various types of phytotoxic compounds that are difficult to artificially synthesize. Such compounds often represent the lead for new plant growth regulator (Scheffer et al. 1983). In the course of our search for new phytotoxic substances from plant pathogenic fungi, we previously reported that a pair of bicyclo 5-15-fused sesterterpene isomers, referred to as terpestacin (Oka et al. 1993) and 11-epiterpestacin (11-ET, **2**), were isolated from the cultures of two distinct fungal strains, *B. sorokiniana* and *B. cynodontis*,

respectively (Lim et al. 1995; Lim et al. 1996; Nihashi et al. 2002). From the culture of *B. sorokiniana*, an acetylated derivative of 11-ET, fusaproliferin (**3**) was also isolated (Santini et al. 1996). In this paper, we report the isolation and structural analysis of another terpestacin-related compound, 11-ETG (**1**), which was found in the culture of *B. sorokiniana*.

Materials and Methods

Fungus and plant

The fungal strain, *B. sorokiniana* NSDR-011 had originally been isolated from a decayed ryegrass leaf with brown spot lesions. Its identification was based on the DNA sequence of Brn1 (Shimizu et al. 1997). Italian ryegrass (*Lolium multiflorum* Lam.) was used for the assay, the seed of which were purchased from Takii Seed company (Kyoto, Japan).

Spectroscopic measurements

Optical rotations were measured on a Jasco model J-5. The IR spectra were determined on a Pye Unicam SP 3200 Infra-Red Spectrophotometer. Electron ionization mass spectra (EI-MS) were obtained on a Hitachi M-80A. The ¹H and ¹³C NMR spectra were determined on a Bruker AC 300 (300 MHz for ¹H and 75 MHz for ¹³C) and/or Bruker ARX 500 (500 MHz for ¹H and 125 MHz for ¹³C). High performance liquid chromatography (HPLC) was performed on Hitachi L-6200 apparatus equipped with an L-4200H UV-Vis detector using a Cosmosil octadecyl silica (ODS) column. Gas chromatography (GC) was performed by a Hitachi G-5000 equipped with an FID detector using a “Quadrex” bonded methyl silicone fused silica capillary column 0.25 mm×25 m.

Isolation and purification of 11-ETG. The fungus was cultured on a PSA medium in Petri dishes (90 mm i.d.) for 14 days in the dark at 25 °C. The agar plates with mycelial body were soaked in acetone, and the acetone extract was filtered and concentrated under reduced pressure at 40 °C. The resulting aqueous solution was re-extracted with ethyl acetate, and the solvent was removed

Chi-Hwan Lim (✉)
E-mail: chlim@cnu.ac.kr

¹Department of Bio-Environmental Chemistry, College of Agricultural Biology & Life Science, Chungnam National University, Taejeon 305-600, Republic of Korea

²Shionogi & Co, LTD, 3-1-1, Futaba-Cho, Toyonaka, Osaka 561-0825, Japan

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

under reduced pressure to obtain a crude extract. This material was subjected to silica gel column chromatography, using a gradient solvent system from *n*-hexane to ethyl acetate, and then to methanol. 11-ETG was detected as a major component in the methanol fraction by HPLC. The compound was further purified by HPLC with a Cosmosil ODS column, UV detector (254 nm), and a mobile phase of MeOH/H₂O (80/20). 11-ETG was obtained as amorphous solid material with a yield of 47 mg from 1,000 agar plates.

Physicochemical data of 11-ETG

[α]_D +26.7° (c 0.83, CHCl₃), ¹³C NMR (125 MHz, methanol-d₄) δ : 208.9 (C18), 150.2 (C17), 147.8, 137.9, 132.4, 129.0, 124.0, 122.1, 100.0 (C1'), 75.7 (C11), 73.8 (C3'), 72.5 (C5'), 72.2 (C2'), 70.4 (C4'), 70.0 (C24), 61.3 (C6'), 48.9 (C15), 48.7 (C1), 40.1, 38.9, 34.8, 34.5, 29.5, 28.8, 23.5, 15.7, 14.3, 14.1, 13.9, 9.0, ¹H NMR (500 MHz, methanol-d₄) δ : 5.42 (1H, m, H-13), 5.33 (1H, m, H-3), 5.17 (1H, m, H-7), 4.78 (1H, d, *J*=3.7 Hz, H-1'), 4.00–3.55 (7H), 3.40–3.25 (2H), 2.90–2.80 (2H), 2.55–2.00 (7H), 2.00–1.60 (5H), 1.64 (6H, s), 1.56 (3H, s), 1.33 (3H, d, *J*=7.0 Hz, C25-methyl), 0.93 (3H, s), EI-MS *m/z*: 564 (M⁺), 402, 384, 137, 81 (base peak), 55.

Acetylation of 11-ETG

11-ETG (17.6 mg) was treated with acetic anhydride (1.5 mL) and pyridine (1.5 mL) at 40 °C for 24 hours. The mixture was poured into a small amount of ice-water to decompose the unreacted acetic anhydride. The solvents were eliminated by vacuum evaporation followed by a stream of N₂ for 5 h. The product was purified by chromatography on a Cosmosil ODS column (20 i.d.×250 L mm) eluted with 80% aq. MeOH to obtain a pure acetyl derivative as pale-yellow oil (14.6 mg).

¹³C NMR (75 MHz, CDCl₃) δ : 206.9 (C18), 170–165 (acetyl carbonyl×6), 144.9 (C16), 138.2 (C4), 132.8 (C12), 132.3 (C8), 130.8 (C13), 124.2 (C7), 121.1 (C3), 95.8 (C1'), 78.7 (C11), 70.7 (C24), 70.5 (C2'), 69.9 (C3'), 68.7 (C4'), 67.5 (C5'), 62.0 (C6'), 49.9 (C15), 49.8 (C1), 40.0 (C5), 39.0 (C2), 34.6 (C23), 34.4 (C9), 28.6 (C12), 27.7 (C14), 23.9 (C6), 23–20 (acetyl methyl×6), 16.0 (C19), 15.7 (C20), 15.4 (C21), 15.1 (C25), 11.1 (C22), ¹H NMR (300 MHz, CDCl₃) δ : 5.53 (1H, br d, *J*=5.9 Hz, H-13), 5.43 (1H, dd, *J*=9.5, 10.1 Hz, H-3'), 5.2 (1H, m, H-11), 5.20–5.05 (2H, H-5, H-7), 5.05 (1H, dd, *J*=9.8, 9.8 Hz, H-4'), 5.00 (1H, d, *J*=3.7 Hz, H-1'), 4.86 (1H, dd, *J*=3.7, 10.3 Hz, H-2'), 4.26 (1H, dd, *J*=4.5, 12.3 Hz, H-6a'), 4.10 (1H, dd, *J*=2.4, 12.3 Hz, H-6b'), 4.00 (1H, ddd, *J*=2.4, 4.4, 10.2 Hz, H-5'), 3.75 (1H, dd, *J*=7.0, 9.3 Hz, H-24a), 3.49 (1H, dd, *J*=7.4, 9.5 Hz, H-24b), 2.90–2.80 (2H, H-15, H-23), 2.50–1.60 (12H), 2.24–1.98 (18H, s, acetyl methyl×6), 1.65 (3H, s, C20-methyl)

Analysis of the hexose moiety of 11-ETG

11-ETG (3.0 mg) was dissolved in 3 N HCl (5 mL). The reaction mixture was maintained with stirring at 75 °C for 4 h. After the

Fig. 1 Structure of 11-ETG (1), 11-epiterpestacin (2), and fusaproliferin (3)

completion of the reaction, the solvent was evaporated in vacuo. The residue was subjected to C18 Sep-Pak cartridge (Waters), eluting with a H₂O to obtain the hexose fraction. Trimethylsilylating (TMS) Reagent (Tokyo Kasei Kogyo Co., 500 μL) was added to the dried hexose fraction, and the reaction mixture was vigorously stirred for a while, then directly subjected to GC analysis. By using the same procedure, TMS derivatives of D-(+)-glucose, D-(–)-fructose, and D-(+)-galactose were prepared for comparison.

In another experiment, 11-ETG (37 mg) was dissolved in 10% HCl-MeOH (5 mL) and heated under reflux for 6 h. The reaction mixture was neutralized with Ag₂CO₃ and the resultant precipitate was filtered off. The methyl glucoside yielded (13 mg) was purified by using a C18 Sep-Pak cartridge, and then acetylated with acetic anhydride and pyridine. After removing the solvent, the syrup obtained (28 mg) was purified by silica gel TLC (*n*-hexane : ether : tetrahydrofuran=33:66:1) to afford methyl α-D-glucopyranoside tetraacetate (6 mg) and its β-isomer (8 mg).

Phytotoxicity assay

A specified amount of the test compound in MeOH was applied to a Petri dish (35 mm in diameter) in which two layers of Whatman No. 2 filter paper were laid. After the solvent was evaporated, one mL of water containing 0.5% Tween 20 (Nakarai Chemicals Ltd., Kyoto, Japan) was added to the dish to dissolve the test compound. Uniformly germinated seeds were selected, with roots ca. 2 mm in length, and placed in the dish (nine seeds per plate) and incubated in a dark condition at 26 °C for 72 h. The inhibitory activity was evaluated by comparing the root length to that of the control.

Results and Discussion

Structure elucidation of 11-ETG

The ¹H-NMR and ¹³C-NMR spectra of 11-ETG were very similar

Fig. 2 NMR chemical shift change of 11-ETG (1) according to acetylation reaction

to those of the 11-ET except the signals assignable to the hexose protons at δ H 4.2–3.3 and at δ H 4.78, and the carbons at δ C 60–75 and δ C 100. The signal at δ C 100 ppm was characteristic of an anomeric carbon of the sugar moiety. The EI-mass spectra of 11-ETG showed a molecular ion at m/z 564, which is larger than 11-ET (M.W. 402) by 162 m.u. of a dehydrated hexose ($C_6H_{12}O_6 - H_2O$). These data suggested that 11-ETG was a monoglycoside of 11-ET.

Acetylation of 11-ETG afforded a hexaacetyl derivative, which was evidenced by its EI-mass spectrum and 1 H-NMR spectrum. EI-mass spectrum showed $[M]^+$ at m/z 816 ($564 + (CH_3CO \times 6) - 6H$) and the 1 H-NMR spectrum showed the presence of six acetyl methyl groups. The presence of hexose moiety was again supported by a series of typical fragment peaks from an acetylated hexose (m/z 331, 169, 109, 43) in the MS. The six alcoholic hydroxy groups in 11-ETG were reasonably assigned to two hydroxyl groups in the 11-ET moiety and four hydroxyl groups in hexose moiety. Furthermore, the methine proton at δ H 5.00 was correlated to the anomeric carbon at 95.8 ppm in the 1 H- 13 C COSY spectrum, and the 1 H- 1 H COSY data showed the correlation from the signal δ H 5.43 to the six proton signals in the range of δ H 4.0–5.0 ppm in order. By acetylation, the chemical shifts of the signals that corresponded to those of H-11 (δ H 4.00) and C-17 (δ C 150.2) in 11-ETG were moved down field (δ H 5.15 and δ C 170.3) respectively (Fig. 2). The presence of glucose moiety in 11-ETG was also supported by the analysis of the proton coupling constants in the NMR spectrum of 11-ETG. The large vicinal coupling constants observed between H-2' and H-3' (10.3 Hz), H-3' and H-4' (9.8 Hz), and H-4' and H-5' (10.0 Hz) indicated that these four protons were arranged with each other in a trans diaxial configuration, which is consistent with that found in glucopyranose. The configuration of the glucoside bond was determined to be alpha by the small coupling constants between

Fig. 3 Italian ryegrass seedlings root growth inhibitory activity by 11-ETG (1), 11-epiterpestacin (2), and fusaproliferin (3)

anomeric proton H-1' and H-2' ($J=3.7$ Hz).

Based on these findings, 11-ETG was concluded to be α -glucopyranoside of 11-ET at the 24-hydroxyl group.

Phytotoxicity assay

11-ETG (1) did not inhibit the root growth of Italian ryegrass seedlings even at the level of 200 ppm, while its aglycone 11-ET (2) completely inhibited root growth at the level of 100 ppm (Fig. 3). The lower hydrophobicity of 11-ETG relative to 11-ET, due to the presence of glucose moiety, is likely to be responsible for the lower phytotoxicity. This is in contrast with the case of fusicoccins, phytotoxins produced by a plant pathogenic fungus *Phomopsis (Fusicoccum) amygdali*. Fusicoccins are structurally related to 11-ETG in that they are terpenoid (diterpene) glucosides, and it has been shown that the glucose moiety, although it is modified with an additional isoprene moiety, in the toxins is important for the expression of toxicity (Ballio et al. 1964, Toyomasu et al. 2007).

The phytotoxicity of 11-ET suggests that this compound plays an important role in the expression of disease symptoms by this fungus, but it is also possible that the toxin also has more or less negative effect on the growth of the producer fungus itself. If so, the production of 11-ETG in the fungus can serve for detoxification. It is unlikely that plants have comparable detoxification mechanism against such a toxin with specific chemical structure. A comparative analysis of the glycosidation activity of terpenoids between fungi and plants may raise an issue of plant pathological interest.

Acknowledgments This work was supported by a grant from Chungnam National University.

References

- Ballio A, Chain EB, De Leo P, Erlanger BF, Mauri M, Tonolo A (1964) Fusicoccin: a new wilting toxin produced by *Fusicoccum amygdali*. *Nature* 203 (4942): 297
- Lim CH, Miyagawa H, Ueno T, Takenaka H, Sung ND (1996) Siccanol: sesterterpene isolated from pathogenic fungus *Drechslera siccanis*. *Agric Chem Biotechnol* 39: 241–244
- Lim CH, Miyagawa H, Ueno T, Takenaka H, Tsurushima T (1995) Isolation and structure elucidation of terpenoid phytotoxins produced by the plant pathogenic fungus *Bipolaris cynodontis*. In abstracts of Papers of 37th Symposium on the Chemistry of Natural Products, Tokushima, Japan, 325–330
- Nihashi Y, Lim CH, Tanaka C, Miyagawa H, Tamio U (2002) Phytotoxic Sesterterpene, 11-Epiterpestacin, from *Bipolaris sorokiniana* NSDR-011. *Biosci Biotechnol Biochem* 66(3): 685–688
- Oka M, Iimura S, Tenmyo O, Yamamoto H, Kawano K, Hu SL, Fukagawa Y, Oka T (1993) Terpestacin, a new syncytium formation inhibitor from *Arthrinium* sp.. *J Antibiot* 46: 367–373
- Santini A, Ritieni A, Fogriano V, Randazzo G, Manniana L, Benedetti E (1996) Structure and absolute stereochemistry of fusaproliferin, a toxic metabolite from *Fusarium proliferatum*. *J Nat Prod* 59: 109–112
- Scheffer RP, Daly JM, Deverall BJ (1983) *Toxins and plant pathogenesis*, Academic Press, New York
- Shimizu K, Tanaka C, Tsuda M (1997) Cloning of *Brnl*, a reductase gene involved in melanin biosynthesis in *Cochliobolus heterostrophus*. *J Gen Appl Microbiol* 43: 145–150
- Toyomasu T, Tsukahara M, Kaneko A, Niida R, Mitsuhashi W, Dairi T, Kato N, Sassa T (2007) Fusicoccins are biosynthesized by an unusual chimera diterpene synthase in fungi. *PNAS* 104 (9): 3084–3088